

THE GENERALIZED VERSCHIEBUNG MAP FOR CURVES OF GENUS 2

BRIAN OSSERMAN

ABSTRACT. Let C be a smooth curve, and $M_r(C)$ the coarse moduli space of vector bundles of rank r and trivial determinant on C . We examine the generalized Verschiebung map $V_r : M_r(C^{(p)}) \dashrightarrow M_r(C)$ induced by pulling back under Frobenius. Our main result is a computation of the degree of V_2 for a general C of genus 2, in characteristic $p > 2$. We also give several general background results on the Verschiebung in an appendix.

1. INTRODUCTION

In this paper we address the degree of the Verschiebung rational map V_2 induced by pullback under Frobenius on the moduli space M_2 of rank 2 vector bundles with trivial determinant on a smooth proper curve, in the case of genus 2, along with some related questions. We work throughout over an algebraically closed field k of characteristic $p > 2$, except where specified.

Aside from the importance of the Verschiebung in the case of Jacobians, and the consequent desire to understand its generalization to higher rank, motivation for understanding the geometry of the Verschiebung map was provided by the close relationship between the Verschiebung map and characteristic- p representations of the fundamental group of C , when the base field k for our curve C is finite (see the introduction to [16]). In particular, A. J. de Jong showed that curves in the moduli space of vector bundles which are fixed under some iterate of the Verschiebung will correspond to characteristic- p representations for which the geometric fundamental group has infinite image, which he conjectures in [4] cannot happen for characteristic- ℓ representations. He further shows that such curves would have to pass through the undefined locus of the Verschiebung. Another motivation comes from the fact that invariants such as the degree of Verschiebung nearly always seem to be given by polynomials in p , with no obvious explanation for why this should be the case. One might hope that examination of enough different cases of this phenomenon would ultimately provide insight into the general situation.

In order to state our result, we fix some terminology:

Definition 1.1. Let C be a smooth, proper curve. A semistable vector bundle $\mathcal{F}$ on $C^{(p)}$ is said to be **Frobenius-unstable** if $F^*\mathcal{F}$ is unstable. Now suppose C has genus 2. Given a Frobenius-unstable vector bundle $\mathcal{F}$ of rank 2 and trivial determinant, we say that $\mathcal{F}$ is a **reduced** point of the Frobenius-unstable locus if the first-order determinant-preserving infinitesimal deformations of $\mathcal{F}$ inject into the first-order infinitesimal deformations of $F^*\mathcal{F}$.

This paper was partially supported by fellowships from the National Science Foundation and Japan Society for the Promotion of Sciences.

We will show:

Theorem 1.2. *Let C be a smooth, proper genus 2 curve over an algebraically closed field k of characteristic $p > 2$, and suppose that the Frobenius-unstable locus for vector bundles of rank 2 and trivial determinant is composed of δ reduced points. Then:*

- (i) *Each undefined point of V_2 may be resolved by a single blowup, and V_2 has degree $p^3 - \delta$;*
- (ii) *The exceptional divisor associated to such an undefined point maps bijectively to $\mathbb{P}\mathrm{Ext}(\mathcal{L}, \mathcal{L}^{-1}) \subset M_2(C)$, where $\mathcal{L}$ is a theta characteristic on C , and specifically is the destabilizing line bundle for $F^*\mathcal{F}$, where $\mathcal{F}$ is the Frobenius-unstable vector bundle associated to the undefined point.*

It follows from results of Mochizuki (see [20], [28]) or of the author (see [27]) that the hypothesis of Theorem 1.2 holds with $\delta = \frac{2}{3}(p^3 - p)$, so we can conclude the following.

Theorem 1.3. *With the notation of Theorem 1.2, if C is general, we have that $\deg V_2 = \frac{p^3+2p}{3}$. We further know that the undefined points of V_2 may each be resolved by a single blowup, with exceptional divisor mapping bijectively to $\mathbb{P}(\mathrm{Ext}(\mathcal{L}, \mathcal{L}^{-1}))$.*

We begin in Section 2 with some straightforward and general results on degrees of rational maps of projective spaces, and their application to V_2 . We next carry out some calculations involving deformation theory of bundles with connection in Section 3. Finally, in Section 4 we establish, following the argument of Langton's properness theorem, the necessary relationship between the locus of p -curvature 0 connections on the unstable bundles $\mathcal{E}$ of [26, Prop. 2.6], the undefined locus of V_2 , and the image of the exceptional divisor if one blows up to resolve V_2 . Appendix A is a compilation of necessary technical background results on V_n , presented in more generality and including arguments suggested by A. J. de Jong and Christian Pauly. Finally, Appendix B develops some slightly non-standard commutative algebra for non-reduced rings which arises in our vector bundle manipulations.

The existing literature on such geometric questions on the Verschiebung is considerably scarcer than on Frobenius-unstable vector bundles. The only other such results in this area were developed recently by Laszlo and Pauly, who gave explicit polynomials defining the Verschiebung in the particular cases of genus 2, rank 2, and characteristics 2 and 3, in [16] and [17]. Lange and Pauly also obtain our formula for the degree of V_2 via a different approach in [14], although their techniques thus far gives only that it is an upper bound in the case of ordinary curves, and not that it is an equality.

The contents of this paper form a portion of the author's 2004 PhD thesis at MIT, under the direction of Johan de Jong.

ACKNOWLEDGEMENTS

I would like to thank Johan de Jong for his tireless and invaluable guidance. I would also like to thank David Sheppard, David Helm, Brian Conrad, Christian Pauly, and Yves Laszlo for their helpful conversations.

2. PRELIMINARIES ON DEGREES

In this section we make some basic observations about degrees of rational maps from projective space to itself, and then apply these to the specific case of the Verschiebung. We remark that unless otherwise specified, in this section the term ‘point’ shall always refer to a closed point. We suppose we are in the following situation.

Situation 2.1. We are given a rational map $f : \mathbb{P}^n \rightarrow \mathbb{P}^n$ which is dominant and defined at all but a finite set of points. We suppose that we are given homogeneous coordinates X_i on $\mathbb{P}^n$, and f is represented by $n + 1$ homogeneous polynomials F_i of degree d in the X_i .

Although special cases of the following proposition are certainly extremely well-known, the general statement, and in particular the possibility of inequality, appears less widely known.

Proposition 2.2. *In the above situation, we have the inequality:*

$$\deg f \leq d^n - \delta,$$

where δ is the total length of the ‘undefined locus’ subscheme E_f of $\mathbb{P}^n$ cut out by the F_i . Moreover, the following are equivalent:

- a) *The above inequality is an equality;*
- b) *E_f is a locally complete intersection;*
- c) *E_f is Gorenstein.*

In particular, we get equality when the length of the points of E_f are all 1 or 2.

Proof. Choose P in $\mathbb{P}^n$, and write $H_1, \dots, H_n$ for n hyperplanes cutting out P . Denote by E_P the (scheme-theoretic) intersection of the $f^*(H_i)$. Now, for a given X_i on the image space, we observe that on $\mathbb{P}_{f^*X_i}^n = \mathbb{P}_{F_i}^n$, we have $f^{-1}(P) \cong E_P$. As i varies, the $\mathbb{P}_{F_i}^n$ will cover everything except E_f . Since the $f^*(H_i)$ are given by polynomials homogeneous and linear in the F_i , there is a closed immersion of E_f into E_P . Hence, we can write E_P as a set as $E_f \cup f^{-1}(P)$, with a closed immersion of the latter into the former. For P general, $f^{-1}(P)$ and hence E_P is 0-dimensional, with length $f^{-1}(P) = \deg f$. Since $\delta := \text{length } E_f$, and E_P is a complete intersection of n hypersurfaces of degree d , Bezout’s theorem yields the desired inequality.

Now, E_f being a locally complete intersection implies that it is Gorenstein, and conversely, since E_f is cut out by $n + 1$ hypersurfaces, if it is Gorenstein it must be a local complete intersection; see [6, Cor 21.19] and [6, p. 542]. Next, to see that a) implies b), we note if we have equality, $E_f \rightarrow E_P$ is an isomorphism onto its image, so since E_P is a complete intersection, E_f is a locally complete intersection.

Finally, we need to show that if E_f is a locally complete intersection, the ideal generated by the $f^*(H_i)$ corresponding to a general point P on the image $\mathbb{P}^n$ is locally equal to the ideal of the F_i at each of the finitely many Q of E_f ; we can therefore check the statement one Q at a time. We fix some $Q \in E_f$, and assume we have fixed a choice of dehomogenization, so that the F_i are actually functions on a neighborhood of Q . We also note that it will be enough to prove the statement for any particular choice of H_i cutting out the given P . We first observe that since E_f is locally a complete intersection, Nakayama’s lemma implies that the defining ideal I_{E_f} of E_f at Q may be generated by n of the $n + 1$ generators F_i . Hence,

we have $F_j = \sum_{i \neq j} a_i F_i$ for some integer j and $a_i \in \mathcal{O}_{\mathbb{P}^n, Q}$; reindexing, we can assume $j = n + 1$. Now, a general point P in the image $\mathbb{P}^n$ may be cut out by H_i of the form $X_i - \lambda_i X_{n+1}$ for $i \leq n$. One then checks directly that for a general choice of the λ_i , F_{n+1} is in the ideal generated by the $f^*(H_i)$. It follows that the other F_i will also be in the ideal generated by the $f^*(H_i)$, giving $E_f \cong E_P$ at Q , as desired. $\square$

For posterity, we observe that the inequality of the preceding proposition need not be an equality:

Example 2.3. Consider the map from $\mathbb{P}^2$ to itself given by $(X, Y, Z) \mapsto (X^3, Y^3, XYZ)$. This is undefined only at $(0, 0, 1)$, where the subscheme E_f has length 5, and is visibly not a local complete intersection. We can check that the map is dominant of degree 3, so as required, we have strict inequality in the preceding proposition.

We now apply these results to examine what we can say about the degree of the Verschiebung map induced by pullback under Frobenius on the moduli space of vector bundles of rank 2 with trivial determinant on a curve C of genus 2. We begin by reviewing the basic facts about the moduli space and Verschiebung, without any hypotheses on the genus.

We will denote by M_2 the moduli space of semistable vector bundles of rank 2 and trivial determinant on C , which will be the only space we will consider here. We will refer to the map induced by pullback under Frobenius on moduli spaces of vector bundles as the **Verschiebung**, since in the case of line bundles this is precisely what it is. We denote by V_2 the particular Verschiebung map on M_2 . V_2 is a dominant rational map, with finitely many undefined points composed of the Frobenius-unstable vector bundles. See Appendix A for the precise technical definitions and proofs of these statements, as well as [12, Thm. 3.2] for the finiteness result.

Now, in our situation of rank 2 bundles on a curve of genus 2, it is a theorem of Narasimhan and Ramanan that $M_2 \cong \mathbb{P}^3$; see [24, Thm. 2, §7], and note that despite the Riemann surface language, the argument goes through unmodified in arbitrary odd characteristic. We also know that the Verschiebung map is given by polynomials of degree p ; see [17, Prop. A.2], and note that $\mathcal{O}(1)$ on our $\mathbb{P}^3$ pulls back to the inverse of the determinant bundle on the moduli stack, by Lemma A.9. Alternatively, one can argue in our case by pulling back to the Jacobians of C and $C^{(p)}$, which map to Kummer surfaces inside $M_2 \cong \mathbb{P}^3$.

Putting this together with Proposition 2.2 yields the following.

Corollary 2.4. *For curves of genus 2, the degree of V_2 is bounded above by p^3 ; or more sharply, $p^3 - \delta$, where δ is the number of points at which V_2 is undefined. If the undefined points are reduced, this upper bound is an equality.*

On the other hand, we also have:

Lemma 2.5. *For curves of genus 2, the degree of V_2 is bounded below by p^2 .*

Proof. For any $\mathcal{L}$ of degree 0 and not 2-torsion, consider the bundle $\mathcal{L} \oplus \mathcal{L}^{-1}$. There are p^2 line bundles of degree 0 mapping to $\mathcal{L}$ under V_2 on the Jacobian, differing from each other by a p -torsion line bundle, and since $\mathcal{L}$ is not 2-torsion, this gives p^2 bundles mapping to $\mathcal{L} \oplus \mathcal{L}^{-1}$. Thus, on an open part of the Kummer surface, each point has at least p^2 preimages, and noting that the points having

positive-dimensional preimage must have codimension at least 1 in the Kummer surface, we conclude the desired lower bound on the degree of V_2 . $\square$

3. DEFORMATION THEORY CALCULATIONS

We begin by reviewing some fundamental facts about representing groups of deformations with cohomology and hypercohomology. Throughout the remainder of this paper, excepting the appendices, we fix the notation:

Notation 3.1. A ‘deformation’ refers to a first order infinitesimal deformation, and ϵ is a square-zero element.

Let C be any smooth curve, with vector bundle $\mathcal{E}$ and connection ∇ . The following propositions are well-known, and may be checked explicitly in terms of Čech cocycles on the U_i .

Proposition 3.2. *The space of deformations of $\mathcal{E}$ is isomorphic to $H^1(C, \mathcal{E}nd(\mathcal{E}))$, and the space of deformations of $\mathcal{E}$ preserving the determinant of $\mathcal{E}$ is isomorphic to $H^1(C, \mathcal{E}nd^0(\mathcal{E}))$.*

Proposition 3.3. *The space of deformations of ∇ over $\mathcal{E}$ (respectively, fixing the determinant of ∇) is isomorphic to $H^0(C, \mathcal{E}nd(\mathcal{E}) \otimes \Omega_C^1)$ (respectively, $H^0(C, \mathcal{E}nd^0(\mathcal{E}) \otimes \Omega_C^1)$).*

In order to analyze transport of ∇ along automorphisms of $\mathcal{E}$, we introduce the sheaf map which is described by the connection $\nabla^{\mathcal{E}nd}$ induced on $\mathcal{E}nd(\mathcal{E})$ by ∇ ; we denote it $d_\nabla : \mathcal{E}nd(\mathcal{E}) \rightarrow \mathcal{E}nd(\mathcal{E}) \otimes \Omega_C^1$, and it is given by $\phi \mapsto \nabla \circ \phi - \phi \circ \nabla$. We then have the following.

Proposition 3.4. *The space of transport equivalence classes of deformations of ∇ over $\mathcal{E}$ is isomorphic to $H^0(C, \mathcal{E}nd(\mathcal{E}) \otimes \Omega_C^1) / d_\nabla(H^0(C, \mathcal{E}nd(\mathcal{E})))$. Furthermore, $d_\nabla|_{\mathcal{E}nd^0(\mathcal{E})}$ takes values in $\mathcal{E}nd^0(\mathcal{E})$, and if $\text{char } k$ is prime to the rank of $\mathcal{E}$, the space of transport equivalence classes of deformations of ∇ over $\mathcal{E}$ with fixed determinant is isomorphic to $H^0(C, \mathcal{E}nd^0(\mathcal{E}) \otimes \Omega_C^1) / d_\nabla(H^0(C, \mathcal{E}nd^0(\mathcal{E})))$.*

Proof. Beyond the usual cocycle-level verifications, the only observation is that since $\text{char } k$ is prime to the rank, and scalar automorphisms fix connections under transport, we have $d_\nabla(H^0(C, \mathcal{E}nd^0(\mathcal{E}))) = d_\nabla(H^0(C, \mathcal{E}nd(\mathcal{E})))$. $\square$

However, parametrizing pairs of deformations of $\mathcal{E}$ together with deformations of ∇ over $\mathcal{E}$ cannot be described naturally with sheaf cohomology, but rather requires sheaf hypercohomology. Since our complex will have only two terms, hypercohomology in our situation is particularly simple. Given $f : \mathcal{F}_1 \rightarrow \mathcal{F}_2$ an element of $\mathbb{H}^1$ is given by a pair of a Čech 1-cocycle of $\mathcal{F}_1$ and a Čech 0-cochain of $\mathcal{F}_2$ which agree under f and the Čech coboundary map.

Notation 3.5. The complex $\mathcal{E}nd^0(\mathcal{E}) \xrightarrow{d_\nabla} \mathcal{E}nd^0(\mathcal{E}) \otimes \Omega_C^1$ will also be denoted by $\mathcal{K}^\bullet$.

Given the explicit description of hypercohomology, the following proposition may also be checked directly in terms of our trivializations.

Proposition 3.6. *The space of transport equivalence classes of deformations of $\mathcal{E}$ together with ∇ (respectively, deformations fixing both determinants) is isomorphic to $\mathbb{H}^1(C, \mathcal{E}nd(\mathcal{E}) \xrightarrow{d_\nabla} \mathcal{E}nd(\mathcal{E}) \otimes \Omega_C^1)$ (respectively, $\mathbb{H}^1(C, \mathcal{E}nd^0(\mathcal{E}) \xrightarrow{d_\nabla} \mathcal{E}nd^0(\mathcal{E}) \otimes \Omega_C^1)$).*

Now, every deformation ∇' of ∇ on $\mathcal{E}$ gives rise to the obvious pair of deformations of $(\mathcal{E}, \nabla)$: namely, $(\mathcal{E}, \nabla')$. Similarly, a pair $(\mathcal{E}', \nabla')$ yields a deformation $\mathcal{E}'$ of $\mathcal{E}$. This gives us induced maps, but we use the standard spectral sequence for the cohomology of a double complex to show that we obtain a short exact sequence, and obtain some additional information.

Proposition 3.7. *Suppose ∇ has p -curvature 0, and the corresponding $\mathcal{F}$ which pulls back under Frobenius to $\mathcal{E}$ is stable. Then d_∇ is injective on global sections, so we can consider $H^0(C, \mathcal{E}nd^0(\mathcal{E}))$ to be a subgroup of $H^0(C, \mathcal{E}nd^0(\mathcal{E}) \otimes \Omega_C^1)$, and we left get a left exact sequence*

$$0 \rightarrow H^0(C, \mathcal{E}nd^0(\mathcal{E}) \otimes \Omega_C^1) / H^0(C, \mathcal{E}nd^0(\mathcal{E})) \rightarrow \mathbb{H}^1(C, \mathcal{K}^\bullet) \rightarrow H^1(C, \mathcal{E}nd^0(\mathcal{E}))$$

where the image on the right is the image under Frobenius pullback of $H^1(C^{(p)}, \mathcal{E}nd^0(\mathcal{F}))$.

Proof. Starting with the Cech double complex associated to our complex, and taking differentials in the vertical direction, we see that the E_1 term is:

$$\begin{array}{ccccccc} & & \vdots & & \vdots & & \\ & & & & & & \\ & & 0 & & 0 & & \\ & & & & & & \\ H^1(C, \mathcal{E}nd^0(\mathcal{E})) & \xrightarrow{d_2^1} & H^1(C, \mathcal{E}nd^0(\mathcal{E}) \otimes \Omega_C^1) & & 0 & & \cdots \\ & & & & & & \\ H^0(C, \mathcal{E}nd^0(\mathcal{E})) & \xrightarrow{d_2^0} & H^0(C, \mathcal{E}nd^0(\mathcal{E}) \otimes \Omega_C^1) & & 0 & & \cdots \end{array}$$

where d_2^i are the maps induced on Cech i -cocycles by d_∇ . We see immediately that the spectral sequence stabilizes at E_2 , and yields a short exact sequence:

$$0 \rightarrow \text{coker } d_2^0 \rightarrow \mathbb{H}^1(C, \mathcal{E}nd^0(\mathcal{E}) \rightarrow \mathcal{E}nd^0(\mathcal{E}) \otimes \Omega_C^1) \rightarrow \ker d_2^1 \rightarrow 0.$$

An element in the kernel of d_2^0 is a trace zero endomorphism of $\mathcal{E}$ which commutes with ∇ . But this is precisely the condition for it to come from a trace zero endomorphism of $\mathcal{F}$ (see, for instance, [13, Thm 5.1]), which must be 0, since $\mathcal{F}$ is stable (this follows almost immediately from the definition of stability and the fact that there are no non-trivial division algebras over an algebraically closed field; see [11, Cor. 1.2.8]).

Similarly, an element in the kernel of d_2^1 comes from $H^1(C^{(p)}, \mathcal{E}nd^0(\mathcal{F}))$, as asserted. $\square$

What is interesting here is that not every deformation $\mathcal{E}'$ of $\mathcal{E}$ arises in this way; indeed, $\mathcal{E}'$ admits a ∇' if and only if it came from some deformation $\mathcal{F}'$ of $\mathcal{F}$. In particular, a given deformation of $\mathcal{E}$ admits a deformation of ∇ if and only if it admits a deformation of ∇ having p -curvature 0.

We also get some information by computing the spectral sequence for the double complex in the other direction. Specifically, we have:

Proposition 3.8. *With the same notation and hypotheses as in Proposition 3.7, we have*

$$H^1(C^{(p)}, \mathcal{E}nd^0(\mathcal{F})) \hookrightarrow \mathbb{H}^1(C, \mathcal{K}^\bullet)$$

The image is described by 1-cocycles of $\mathcal{E}nd^0(\mathcal{E})$ in the kernel of d_∇ (together with the zero 0-cochain of $\mathcal{E}nd^0(\mathcal{E}) \otimes \Omega_C^1$).

Proof. Taking differentials in the opposite direction as before, we find our E_2 term looks like

$$\begin{array}{ccccccc}
 & \vdots & & \vdots & & \vdots & \\
 & & & & & & \\
 H^2(C, \ker d_\nabla) & & H^2(C, \operatorname{coker} d_\nabla) & & 0 & & \cdots \\
 & & & & & & \\
 H^1(C, \ker d_\nabla) & & H^1(C, \operatorname{coker} d_\nabla) & & 0 & & \cdots \\
 & & & & & & \\
 H^0(C, \ker d_\nabla) & & H^0(C, \operatorname{coker} d_\nabla) & & 0 & & \cdots
 \end{array}$$

Now, $H^1(C, \ker d_\nabla)$ cannot have any further nonzero differentials mapping into or out of it, so it injects into $\mathbb{H}^1(C, \mathcal{K}^\bullet)$. We saw in the proof of the previous proposition that $\ker d_\nabla$ is precisely $F^{-1}\mathcal{E}nd^0(\mathcal{F})$. Since F is a homeomorphism, we have $H^1(C, \ker d_\nabla) = H^1(C^{(p)}, \mathcal{E}nd^0(\mathcal{F}))$, and we thus obtain the desired injection, and description of its image. $\square$

The injectivity may be seen equally easily from the categorical equivalence between vector bundles $\mathcal{F}$ on $C^{(p)}$ and pairs $(\mathcal{E}, \nabla)$ on C . Note, however, that this proposition does not imply that every non-trivial deformation of $\mathcal{F}$ maps to a non-trivial deformation of $\mathcal{E}$.

We now suppose that $\mathcal{E}$ is an extension of $\mathcal{L}^{-1}$ by $\mathcal{L}$, with $\mathcal{L}$ any line bundle on C . The short exact sequence

$$0 \longrightarrow \mathcal{L} \xrightarrow{i} \mathcal{E} \xrightarrow{j} \mathcal{L}^{-1} \longrightarrow 0$$

induces maps

$$(3.1) \quad d_1 : \operatorname{Hom}(\mathcal{L}^{-1}, \mathcal{L}) \rightarrow \mathcal{E}nd^0(\mathcal{E})$$

and

$$(3.2) \quad d_2 : \mathcal{E}nd^0(\mathcal{E}) \rightarrow \operatorname{Hom}(\mathcal{L}, \mathcal{L}^{-1})$$

by composition. Specifically, $d_1(\phi) = i \circ \phi \circ j$, and $d_2(\phi) = j \circ \phi \circ i$. Thus, we obtain natural candidates for maps $H^0(C, \operatorname{Hom}(\mathcal{L}^{-1}, \mathcal{L}) \otimes \Omega_C^1) \rightarrow \mathbb{H}^1(C, \mathcal{K}^\bullet)$ and $\mathbb{H}^1(C, \mathcal{K}^\bullet) \rightarrow H^1(C, \operatorname{Hom}(\mathcal{L}, \mathcal{L}^{-1}))$, although we will need to check that the first is well-defined. We now restrict to our particular situation of interest, for the remainder of this section and the next.

Situation 3.9. Suppose that C has genus 2, and $\mathcal{E}$ an extension of $\mathcal{L}^{-1}$ by $\mathcal{L}$, with $\mathcal{L}$ a theta characteristic. Suppose also that the connection ∇ on $\mathcal{E}$ has vanishing p -curvature.

Proposition 3.10. *In our situation, the maps d_1 and d_2 induce a short exact sequence*

$$0 \rightarrow \Gamma(\operatorname{Hom}(\mathcal{L}^{-1}, \mathcal{L}) \otimes \Omega_C^1) \rightarrow \mathbb{H}^1(C, \mathcal{K}^\bullet) \rightarrow \operatorname{Ext}(\mathcal{L}, \mathcal{L}^{-1}) \rightarrow 0$$

Proof. It is convenient to introduce the filtration of $\mathcal{E}nd^0(\mathcal{E})$ given by $0 \subset \text{im } d_1 \subset \ker d_2 \subset \mathcal{E}nd^0(\mathcal{E})$. We claim this induces the following filtration of our original complex:

$$(3.3) \quad \begin{array}{ccc} \mathcal{E}nd^0(\mathcal{E}) & \xrightarrow{d_\nabla} & \mathcal{E}nd^0(\mathcal{E}) \otimes \Omega_C^1 \\ \uparrow & & \uparrow \\ \ker d_2 & \xrightarrow{d_\nabla} & \mathcal{E}nd^0(\mathcal{E}) \otimes \Omega_C^1 \\ \uparrow & & \uparrow \\ \text{im } d_1 & \xrightarrow{d_\nabla} & (\ker d_2) \otimes \Omega_C^1 \\ \uparrow & & \uparrow \\ 0 & \xrightarrow{d_\nabla} & (\text{im } d_1) \otimes \Omega_C^1 \end{array}$$

The only part which requires any verification is that $d_\nabla(\text{im } d_1) \subset (\ker d_2) \otimes \Omega_C^1$, which is easily checked directly. Given this filtration of our complex, we get (see [5, 1.4.5]) a spectral sequence converging to its hypercohomology, whose E_2 term is given in terms of the hypercohomology of the quotient complexes. Specifically, if $\mathcal{K}^\bullet$ is our complex, and $\mathcal{A}_i^\bullet$ the filtration, we get $E_2^{p,q} = \mathbb{H}^{p+q}(C, \text{Gr}_{\mathcal{A}}^q(\mathcal{K})) \Rightarrow \mathbb{H}^{p+q}(C, \mathcal{K})$. We must therefore start by calculating the associated graded complexes of the filtration. Now, we have $\text{im } d_1 \cong \mathcal{H}om(\mathcal{L}^{-1}, \mathcal{L})$, $\ker d_2 / \text{im } d_1 \cong \mathcal{O}_C$, and $\mathcal{E}nd^0(E) / \ker d_2 \cong \mathcal{H}om(\mathcal{L}, \mathcal{L}^{-1})$. Of course, the associated graded sheaves on the right are gotten by taking these, shifted them down by one, and tensoring with Ω_C^1 . Noting that $\mathcal{H}om(\mathcal{L}^{-1}, \mathcal{L}) \cong \mathcal{L}^{\otimes 2} \cong \Omega_C^1$, we actually get isomorphic line bundles for the middle two associated graded complexes, and we have to check that the maps between them are isomorphisms. One can carry this out directly in terms of transition matrices, by checking that the two maps are linear and non-zero, at least in odd characteristic.

Noting that the hypercohomology of an isomorphism vanishes, and the hypercohomology of a 1-term complex concentrated in the i th place is the cohomology of the nonzero term, shifted by i , we obtain the E_2 term of our spectral sequence:

$$\begin{array}{ccccccc} H^0(C, \mathcal{H}) & H^1(C, \mathcal{H}) & 0 & \cdots & & & \\ & 0 & 0 & 0 & \cdots & & \\ & & 0 & 0 & 0 & \cdots & \\ & & & 0 & H^0(C, \mathcal{H} \otimes \Omega_C^1) & H^1(C, \mathcal{H} \otimes \Omega_C^1) & \end{array}$$

where $\mathcal{H} := \mathcal{H}om(\mathcal{L}, \mathcal{L}^{-1})$.

Now, there are potentially non-zero differentials between the remaining terms, but because $H^0(C, \mathcal{H}om(\mathcal{L}, \mathcal{L}^{-1})) = 0$, the $H^0(C, \mathcal{H} \otimes \Omega_C^1)$ cannot have any further nonzero differentials, and applying $H^1(C, \mathcal{H}om(\mathcal{L}, \mathcal{L}^{-1})) = \text{Ext}(\mathcal{L}, \mathcal{L}^{-1})$, we obtain the desired short exact sequence for $\mathbb{H}^1(C, \mathcal{K}^\bullet)$. $\square$

We also note from the construction of the spectral sequence that the map from $\mathbb{H}^1(C, \mathcal{K}^\bullet)$ to $\text{Ext}(\mathcal{L}, \mathcal{L}^{-1})$ is, at least up to sign, the map induced by first mapping to $H^1(C, \mathcal{E}nd^0(\mathcal{E}))$, and then taking the map induced by d_2 on H^1 . We thus get a diagram:

$$\begin{array}{ccccc}
 & & H^0(C, \mathcal{E}nd^0(\mathcal{E}) \otimes \Omega_C^1) / H^0(C, \mathcal{E}nd^0(\mathcal{E})) & & \\
 & \swarrow & \downarrow & & \\
 H^0(C, (\Omega_C^1)^2) & \xrightarrow{\quad} & \mathbb{H}^1(C, \mathcal{K}^\bullet) & \xrightarrow{\quad} & \text{Ext}(\mathcal{L}, \mathcal{L}^{-1}) \\
 & & \downarrow & \nearrow & \\
 & & H^1(C, \mathcal{E}nd^0(\mathcal{E})) & &
 \end{array}$$

where the inclusion on the upper left follows formally from exactness in the middle. But now we compare the dimensions to conclude:

Proposition 3.11. *The kernel of $\mathbb{H}^1(C, \mathcal{K}^\bullet) \rightarrow H^1(C, \mathcal{E}nd^0(\mathcal{E}))$ is equal to the kernel of $\mathbb{H}^1(C, \mathcal{K}^\bullet) \rightarrow \text{Ext}(\mathcal{L}, \mathcal{L}^{-1})$. Equivalently, a deformation of the pair $(\mathcal{E}, \nabla)$ induces the trivial extension in $\text{Ext}(\mathcal{L}, \mathcal{L}^{-1})$ if and only if it was the trivial deformation of $\mathcal{E}$ (together with any deformation of ∇).*

Proof. We need only show that the dimensions of the two kernels are equal. Applying the Riemann-Roch theorem for vector bundles together with the self-duality of $\mathcal{E}nd^0(\mathcal{E})$, we compute that both

$$h^0(C, (\Omega_C^1)^{\otimes 2}) = h^0(C, \mathcal{E}nd^0(\mathcal{E}) \otimes \Omega_C^1) - h^0(C, \mathcal{E}nd^0(\mathcal{E})) = 3,$$

as desired. $\square$

4. GEOMETRIC SIGNIFICANCE

In this section, we lend some geometric substance to what we have calculated thus far, largely by characterizing the map $H^1(C, \mathcal{E}nd^0(\mathcal{E})) \rightarrow \text{Ext}(\mathcal{L}, \mathcal{L}^{-1})$ in geometric terms. We continue with the hypotheses of Situation 3.9. Suppose we have a family of vector bundles $\tilde{\mathcal{E}}$ on C with base T , and trivial determinant, such that for some k -valued point $0 \in T$, $\tilde{\mathcal{E}}|_0 \cong \mathcal{E}$. Since we have a map $\mathcal{E} \rightarrow \mathcal{L}^{-1}$, by adjointness we get a map $\tilde{\mathcal{E}} \rightarrow i_{0*}\mathcal{L}^{-1}$ (see Lemma B.11), and we can take the kernel to get a new family $\tilde{\mathcal{E}}'$ over T which is isomorphic to $\tilde{\mathcal{E}}$ away from 0 (this will be another vector bundle if T is a reduced curve, but not quite, for instance, if T is $\text{Spec } k[\epsilon]$; see Theorem B.10 of the appendix).

We can then restrict back to the fiber at 0, where we will get a new $\mathcal{E}'$ on C which will also be a vector bundle (even if $T = \text{Spec } k[\epsilon]$), of rank 2 and trivial determinant. Everything but the trivial determinant assertion actually follows immediately from Theorem B.10, with T either a curve or $\text{Spec } k[\epsilon]$. For the triviality of the determinant, and consequent remarks, we will for the moment assume that T is a curve. In this case, we get the desired result from [22, Cor 5.6], since we have triviality of the determinant away from 0 on T .

Moreover, we see that $\mathcal{E}'$ is an extension of $\mathcal{L}$ by $\mathcal{L}^{-1}$: Since i is a closed immersion, $(i_{0*}\mathcal{L}^{-1})|_0 = \mathcal{L}^{-1}$, so we have an exact sequence

$$(4.1) \quad \mathcal{E}' \rightarrow \mathcal{E} \rightarrow \mathcal{L}^{-1} \rightarrow 0$$

but since the kernel of $\mathcal{E} \rightarrow \mathcal{L}^{-1}$ is $\mathcal{L}$, $\mathcal{E}' \rightarrow \mathcal{E}$ factors through $\mathcal{L} \rightarrow \mathcal{E}$, and we see that we get a map from $\mathcal{E}'$ to $\mathcal{L}$, which is surjective, by exactness of equation 4.1. Since $\mathcal{E}'$ has trivial determinant, it follows that $\mathcal{E}'$ is an extension of $\mathcal{L}$ by $\mathcal{L}^{-1}$. Thus, in the case that T is a curve, we get a map

$$\phi_{\mathcal{E}} : \{\tilde{\mathcal{E}} \text{ over } T\} \rightarrow \text{Ext}(\mathcal{L}, \mathcal{L}^{-1}),$$

This will have the following significance for us in trying to understand the Verschiebung:

Lemma 4.1. *Suppose T is a curve, 0 a point of T , and $\tilde{\mathcal{F}}$ a nontrivial family of semistable vector bundles over T with trivial determinant, such that $F^*\tilde{\mathcal{F}} = \mathcal{E}$, where $\mathcal{F} := \tilde{\mathcal{F}}|_0$. That is to say, $\tilde{\mathcal{F}}$ gives a nonconstant map of T into the moduli space M_2 , passing through a point where V_2 is undefined. Then writing $\mathcal{E} = F^*\tilde{\mathcal{F}}$, if $\mathcal{E}' = \phi_{\mathcal{E}}(\mathcal{E}) \neq 0$, the limit point of the image of T under V_2 at $\mathcal{F}$ is given by $\mathcal{E}'$.*

Proof. Since $\tilde{\mathcal{E}}'$ is isomorphic to $\tilde{\mathcal{E}}$ away from the limit point, it suffices to check that $\mathcal{E}'$ is semistable as long as it is nonzero in $\text{Ext}(\mathcal{L}, \mathcal{L}^{-1})$. But this may be checked directly from the definitions, using the fact that $\mathcal{L}$ has degree 1. $\square$

The implication is that if we are lucky, we will be able to describe the image of the exceptional divisor of V_2 in terms of $\mathbb{P}\text{Ext}(\mathcal{L}, \mathcal{L}^{-1})$, which can easily be checked to define a hyperplane inside of M_2 , using the identification $M_2 \cong \mathbb{P}H^0(\text{Pic}^1(C), \mathcal{O}(2\Theta))$, together with [24, Lem. 5.8] to handle the semi-stable locus. To show that this hyperplane will in fact be the image of the exceptional divisor, we first note that if $T = \text{Spec } k[\epsilon]$, we actually still get a map $\phi_{\mathcal{E}} : \text{Def}^0(\mathcal{E}) \rightarrow \text{Ext}(\mathcal{L}, \mathcal{L}^{-1})$. In fact, more specifically, we have:

Lemma 4.2. *$\phi_{\mathcal{E}}$ still exists in the case $T = \text{Spec } k[\epsilon]$, inducing a map from $\text{Def}^0(\mathcal{E})$ to $\text{Ext}(\mathcal{L}, \mathcal{L}^{-1})$ which arises as the negative of the map on 1-cocycles induced by $d_2 : \text{End}^0(\mathcal{E}) \rightarrow \text{Hom}(\mathcal{L}, \mathcal{L}^{-1})$ (3.2), from the previous section.*

Proof. Let U_1, U_2 be an open cover of C trivializing $\mathcal{L}$, $\mathcal{E}$, and Ω_C^1 , and ω_i one-forms trivializing Ω_C^1 on the U_i . In addition, we set the convention that all 1-cocycles will be written with coordinates on U_2 . To prove the lemma, we have to start by pinning down the identification of $\text{Ext}(\mathcal{L}, \mathcal{L}^{-1})$ with $H^1(C, \text{Hom}(\mathcal{L}, \mathcal{L}^{-1}))$. We will think of $\text{Ext}(\mathcal{L}, \mathcal{L}^{-1})$ as being described by transition matrices of the form $F = \begin{bmatrix} \varphi_{12} & 0 \\ f & \varphi_{12}^{-1} \end{bmatrix}$, where f is any regular section on $U_1 \cap U_2$. We first claim that rather than taking f itself as our 1-cocycle of $\text{Hom}(\mathcal{L}, \mathcal{L}^{-1})$, we will have to take $-\varphi_{12}^{-1}f$. Indeed, this may be checked directly by following through the standard identification: starting from an extension $\mathcal{F} \in \text{Ext}(\mathcal{L}, \mathcal{L}^{-1})$, the corresponding 1-cocycle of $\text{Hom}(\mathcal{L}, \mathcal{L}^{-1})$ is obtained by considering the exact sequence

$$0 \rightarrow \text{Hom}(\mathcal{L}, \mathcal{L}^{-1}) \rightarrow \text{Hom}(\mathcal{L}, \mathcal{F}) \rightarrow \text{Hom}(\mathcal{L}, \mathcal{L}) \rightarrow 0,$$

and looking at the image in $H^1(C, \text{Hom}(\mathcal{L}, \mathcal{L}^{-1}))$ of the identity in $\text{Hom}(\mathcal{L}, \mathcal{L})$ under the boundary map.

Next, suppose we have a deformation $\mathcal{E}'$ of $\mathcal{E}$ given by the transition matrix

$$E(I + \epsilon E') = \begin{bmatrix} \varphi_{12} + \epsilon e_{11} & \varphi_{12}^{-2} + \epsilon e_{12} \\ \epsilon e_{21} & \varphi_{12}^{-1} + \epsilon e_{22} \end{bmatrix}$$

Recalling that d_2 , by definition, took the lower left coordinate of a matrix, and noting that the lower left coordinate of E' will be $\varphi_{12}e_{21}$, we see from the above that it suffices to show that $\phi_{\mathcal{E}}(E')$ is described by the transition matrix $\begin{bmatrix} \varphi_{12} & 0 \\ e_{21} & \varphi_{12}^{-1} \end{bmatrix}$. But we can calculate $\phi_{\mathcal{E}}(E')$ directly, in this case. Denote by s_i, t_i our trivializing basis on U_i , in terms of which E and E' are written. Also write u_i for the trivialization of $\mathcal{L}^{-1}$ on U_i . This means we have $s_2 = (\varphi_{12} + \epsilon e_{11})s_1 + \epsilon e_{21}t_1$, $t_2 = (\varphi_{12}^{-2} + \epsilon e_{12})s_1 + (\varphi_{12}^{-1} + \epsilon e_{22})t_1$, and $u_2 = \varphi_{12}^{-1}u_1$. Now, the induced map from $\mathcal{E}'$ to $\mathcal{L}^{-1}$ sends $as_i + bs_i + ct_i + det_i$ simply to cu_i . This means that its kernel is generated by s_i, et_i on U_i . Using the above formulas for s_2, t_2 in terms of s_1, t_1 , we find that this kernel has transition matrix

$$\begin{bmatrix} \varphi_{12} + \epsilon e_{11} & \epsilon \varphi_{12}^{-2} \\ e_{21} & \varphi_{12}^{-1} \end{bmatrix}.$$

Upon modding out by ϵ , this gives precisely the desired form for $\phi_{\mathcal{E}}(E')$. The theory developed in the appendix, and particularly Corollary B.3 and the subsequent discussion, justifies this calculation, even though the transition matrix is not unique, and the kernel itself (prior to restriction) is not characterized by it. $\square$

We next address some compatibility statements on curves and deformations.

Lemma 4.3. *In the same situation as Lemma 4.1, if we write $\tilde{\mathcal{F}}$ for the induced first-order deformation of $\mathcal{F}$ gotten via some closed immersion $t_\epsilon : \text{Spec } k[\epsilon] \hookrightarrow T$ deforming the point $0 \in T$, and $\bar{\mathcal{E}} := F^*\tilde{\mathcal{F}}$, we have $\bar{\mathcal{E}} \cong t_\epsilon^*\tilde{\mathcal{E}}$. Further, $\phi_{\mathcal{E}}\bar{\mathcal{E}} = \phi_{\mathcal{E}}\tilde{\mathcal{E}}$.*

Proof. The first statement follows immediately from the fact that $\tilde{\mathcal{F}}$ is a sheaf on $C^{(p)} \times T$, and all we are saying is that pullback under t_ϵ commutes with pullback under F . Since t_ϵ only acts on T , and F only acts on $C^{(p)}$, they commute.

The second half of the lemma is just an application of Theorem B.12 in the appendix to our specific situation. $\square$

Putting together this lemma with Lemma 4.1, we see:

Theorem 4.4. *In the same situation as the previous lemma, if we suppose $\phi_{\mathcal{E}} \circ F^*$ is injective on first-order deformations of $\mathcal{F}$, then the limit point at 0 of the image of $T \rightarrow M_2$ under the Verschiebung is given as $\phi_{\mathcal{E}} \circ F^*(\mathcal{F})$. In particular, all such limit points are contained in $\mathbb{P}\text{Ext}(\mathcal{L}, \mathcal{L}^{-1}) \subset M_2$. Further, the Verschiebung only needs to be blown up once at $\mathcal{F}$.*

Proof. Everything but the last assertion follows directly from the two lemmas. To show that the Verschiebung only needs to be blown up once at $\mathcal{F}$, it suffices to know that every smooth curve through $\mathcal{F}$ has the limit point of its image under V_2 determined by its tangent at $\mathcal{F}$. It suffices to show that every such curve has a corresponding family of vector bundles on it, since we could then apply our result that it suffices to look at the first order deformation induced by the family. However, because all of our $\mathcal{F}$'s are in the stable locus of our moduli space, the obstruction to a universal bundle is given by a Brauer class; see [11, Cor. 4.3.5], [21, Cor. 2.5, p. 55, Prop. 0.9, p. 16], and [10, Thm. 11.7] to conclude that the GIT quotient is an étale principal PGL_n -bundle even in positive characteristic, and then apply [3, Prop. 3.3.2]. Because k is algebraically closed, by Tsen's theorem [8, Rem. 1.14], [9, Cor. 1.10] we obtain the required bundles on smooth curves. $\square$

Finally, we can draw some conclusions which have immediate consequences for our understanding of the Verschiebung:

Theorem 4.5. *Given $\mathcal{F}$ such that $F^*\mathcal{F} \cong \mathcal{E}$, let $E_{\mathcal{F}}$ be the exceptional divisor above $\mathcal{F}$ after blowing up M_2 to make the Verschiebung a morphism. Then of the following, a) and b) are equivalent, and either implies c) and d):*

- a) *The scheme of connections with vanishing p -curvature on $\mathcal{E}$ is reduced at the point corresponding to $\mathcal{F}$.*
- b) *The map $\text{Def}(\mathcal{F}) \rightarrow \text{Def}(\mathcal{E})$ induced by F^* is injective.*
- c) *The image of $E_{\mathcal{F}}$ under V_2 in M_2 is precisely $\mathbb{P}\text{Ext}(\mathcal{L}, \mathcal{L}^{-1})$.*
- d) *V_2 only needs to be blown up once at $\mathcal{F}$*

Proof. a) is equivalent to there not being any non-trivial deformations of ∇ which hold $\mathcal{E}$ fixed, and have p -curvature 0. The equivalence of this with the condition that there are no nontrivial deformations of $\mathcal{F}$ which pull back to the trivial deformation of $\mathcal{E}$ follows from the categorical equivalence provided by the Cartier isomorphism; see [13, Thm. 5.1].

On the other hand, b) is equivalent to the map $\text{Def}(\mathcal{F}) \rightarrow \text{Ext}(\mathcal{L}, \mathcal{L}^{-1})$ being injective, by Proposition 3.11, which implies it is an isomorphism, since both spaces have dimension 3 over k . Noting that Lemma 4.2 tells us our geometric and cocycles versions of this map are really the same up to sign, by the preceding theorem this implies c), as desired.

Lastly, the fact that d) follows from these conditions also follows from the preceding theorem, since as we just noted, b) gives us that $\text{Def}(\mathcal{F}) \hookrightarrow \text{Ext}(\mathcal{L}, \mathcal{L}^{-1})$. $\square$

We may now easily put everything together to prove our main theorem:

Proof of Theorem 1.2. The assertion of (i) that each Frobenius-unstable bundle corresponds to an undefined point will follow from (iii), since the image of the exceptional divisor of a blow-up centered at such a point is not just a single point. The degree statement then follows from (ii) by Corollary 2.4. But now (ii) and (iii) follow from the implications b) implies c) and d) of Theorem 4.5. $\square$

We conclude with some further questions:

Question 4.6. Are statements c) and d) of Theorem 4.5 in fact equivalent to a) and b)?

Question 4.7. Is the scheme of Frobenius-unstable bundles of rank two and trivial determinant (more precisely, the scheme of transport-equivalence classes of connections with trivial determinant and vanishing p -curvature on the appropriate unstable bundles) isomorphic to the scheme-theoretic undefined locus of V_2 ?

Question 4.8. Is the degree of V_2 constant over all smooth curves of genus 2?

We remark that an affirmative answer to the second question would give an affirmative answer to the third, thanks to a result of Mochizuki [20, II, Thm. 2.8, p. 153]: this gives that our scheme of Frobenius-unstable bundles is finite flat over our space of curves, and that it is smooth over the base field, from which it follows that its fiber over any fixed curve is a local complete intersection.

More generally, one could ask:

Question 4.9. How might one attempt to compute the degree of the Verschiebung for curves of higher genus, or vector bundles of higher rank?

We remark finally that to attempt to address this last question via similar techniques to those of this chapter, it would be necessary not only to generalize our understanding of the undefined locus of the Verschiebung, but also to appropriately generalize the degree formula of Proposition 2.2 to a substantially more general class of projective varieties; indeed, for higher genus and rank, the moduli spaces in question become singular along the strictly semi-stable locus.

APPENDIX A. SOME GENERAL RESULTS ON THE VERSCHIEBUNG

This appendix consists of the formal construction of, and some general results on, the generalized Verschiebung map on coarse moduli spaces of vector bundles. Most of the arguments are due to A. J. de Jong, or Christian Pauly, as indicated. We will work in the situation:

Situation A.1. C is a smooth, proper curve over an algebraically closed field k of characteristic p . $C^{(p)}$ is the p -twist of C over k , and F is the relative Frobenius morphism from C to $C^{(p)}$.

Note that in this appendix, we do not assume $p > 2$.

Remark A.2. The algebraically closed hypothesis is not actually necessary; the moduli space construction can be made to work over a non-algebraically closed field, and the arguments here will go through in this setting. However, it is harder to find references for the general case, and we will only apply the results here in the case of an algebraically closed base field.

Remark A.3. There are a few points to be careful of with respect to characteristic and the general theory of moduli of vector bundles. The main obstruction is boundedness, which is not a problem in our situation of the base being a curve (see [11, Cor. 1.7.7]), and is now known in any dimension via the more involved argument of [15]. However, in characteristic p the statement that the moduli space universally corepresents the relevant functor is in fact no longer true. It is however true that it **uniformly** corepresents the functor, which is to say that it is universal for flat base change, and this is all we will need; see [21, Thm. 1.10, p. 38].

We recall the following definition and theorem:

Definition A.4. Two vector bundles $\mathcal{E}, \mathcal{E}'$ of degree 0 are **S -equivalent** if and only if there are filtrations $F_{\mathcal{E}}$ and $F_{\mathcal{E}'}$ with $\mathrm{Gr}_{F_{\mathcal{E}}} \cong \mathrm{Gr}_{F_{\mathcal{E}'}}$ (this isomorphism is not required to preserve the grading), and the quotients of $F_{\mathcal{E}}$ and $F_{\mathcal{E}'}$ all stable sheaves of degree 0.

Theorem A.5. [11, Thm. 4.3.4] *There is a coarse moduli scheme $M_r(C)$ which uniformly corepresents the functor of semistable vector bundles on C of rank r and trivial determinant. The closed points of $M_r(C)$ correspond to S -equivalence classes of vector bundles; in particular, there is an open subscheme $M_r^s(C)$ whose closed points parametrize stable vector bundles.*

The main results on the Verschiebung are the following. The main arguments for existence and dominance are drawn from unpublished work of de Jong, while Pauly suggested the first portion of the argument for showing that the map doesn't extend over the locus of Frobenius-unstable bundles.

Theorem A.6. *In the above situation, and given $r > 0$, the operation of pulling back vector bundles under F induces a **generalized Verschiebung** rational map $V_r : M_r(C^{(p)}) \dashrightarrow M_r(C)$. If we denote by U_r the open subset of $M_r(C^{(p)})$ corresponding to bundles $\mathcal{E}$ such that $F^*(\mathcal{E})$ is semi-stable, we have further:*

- (i) *The domain of definition of V_r is precisely U_r ;*
- (ii) *V_r is dominant.*

The existence portion of the theorem is straightforward.

Proposition A.7. *The V_r of Theorem A.6 exists, and induces a morphism on U_r .*

Proof. Since semistability is an open condition (see [11, Prop. 2.3.1]), we get an open subfunctor $\mathcal{U}_r$ of the moduli functor $\mathcal{M}_r(C^{(p)})$ corresponding to semi-stable vector bundles on $C^{(p)}$ which pull back under F to semi-stable vector bundles on C . We claim that it is enough to show that this subfunctor is stable under S -equivalence. Indeed, given this, and using the description of the closed points of $M_r(C^{(p)})$ and the fact that it uniformly corepresents $\mathcal{M}_r(C^{(p)})$, it is straightforward to check that $\mathcal{U}_r$ corresponds naturally to an open subscheme U_r of $M_r(C^{(p)})$ which corepresents $\mathcal{U}_r$ and whose closed points are precisely S -equivalence classes of vector bundles on $C^{(p)}$ whose pullbacks under F are semi-stable. Now, Frobenius pullback induces a map from $\mathcal{U}_r$ to $\mathcal{M}_r(C)$; if we compose with the map $\mathcal{M}_r(C) \rightarrow M_r(C)$, since U_r corepresents $\mathcal{U}_r$, we obtain the desired morphism $V_r : U_r \rightarrow M_r(C)$.

We therefore show that our subfunctor is in fact stable under S -equivalence. Let $\mathcal{E}, \mathcal{E}'$ be as in Definition A.4. Since C is smooth, we know F is flat, so F^* behaves well with respect to filtrations and the operation Gr , and we claim this implies that $F^*\mathcal{E}$ is semi-stable if and only if $F^*\text{Gr}_{F_{\mathcal{E}}}$ is semi-stable. Certainly, if $\mathcal{F} \subset F^*\mathcal{E}$ is a destabilizing subsheaf, then by considering the smallest subsheaf in the filtration $F^*F_{\mathcal{E}}$ into which $\mathcal{F}$ maps, there will be a nonzero map of $\mathcal{F}$ into the corresponding quotient in $F^*\text{Gr}_{F_{\mathcal{E}}}$. Conversely, suppose that $\mathcal{F}$ is a destabilizing subsheaf of some quotient in $F^*\text{Gr}_{F_{\mathcal{E}}}$; that is, $\mathcal{F}$ has positive degree and there is an injection into $F^*F_{\mathcal{E}}^{(i)} / F^*F_{\mathcal{E}}^{(i+1)}$ for some i , from which it follows that the cokernel has negative degree. If $\mathcal{F}'$ is the subsheaf of $F^*F_{\mathcal{E}}^{(i)}$ generated by $\mathcal{F}$ and $F^*F_{\mathcal{E}}^{(i+1)}$, the cokernel of the inclusion is the same, hence has negative degree, and it follows that $\mathcal{F}'$ has positive degree and maps into $F^*\mathcal{E}$, giving the desired instability. We conclude that $F^*\mathcal{E}$ is semi-stable if and only if $F^*\mathcal{E}'$ is semi-stable, as desired. $\square$

Before completing the proof of the theorem, we recall a number of well-known facts about moduli spaces of vector bundles.

Theorem A.8. *Let $\mathcal{M}_r(C)$ and $\mathcal{M}_r^{ss}(C)$ denote the Artin stacks of vector bundles of rank r and trivial determinant on C and the open substack of semistable bundles respectively. We have:*

- (i) *The Picard groups of $\mathcal{M}_r(C)$ and $\mathcal{M}_r^{ss}(C)$ are both isomorphic to $\mathbb{Z}$, and generated by the inverse of the determinant line bundle, which we denote by $\mathcal{L}$.*
- (ii) *For n sufficiently large, $\mathcal{L}^n$ has base locus precisely equal to the locus of unstable bundles, and descends to a very ample line bundle on $M_r(C)$.*
- (iii) *$M_r(C)$ is normal, and $\mathbb{Q}$ -factorial.*

Proof. To see that $\text{Pic}(\mathcal{M}_r(C)) \cong \text{Pic}(\mathcal{M}_r^{ss}(C))$ under the natural map, use the argument of [2, Prop. 8.3]. Faltings gives a characteristic-independent argument for the isomorphism with $\mathbb{Z}$ in [7, Thm. 17].

The statement of (ii) in the context of GIT is essentially standard. Specifically, let $\bar{R}$ be a GIT rigidification space whose quotient is $M_r(C)$ [11, §4.3], with linearized bundle $\hat{\mathcal{L}}$, and we will denote by $\hat{R}$ the open subscheme (containing R) of locally free sheaves which are globally generated with vanishing H^1 , after twisting by the power of the ample bundle on C used in the chosen rigidification. We see from vanishing of obstructions on curves and the vanishing H^1 condition that $\hat{R}$ is smooth. The GIT-semistable points of $\hat{R}$ are precisely the semistable bundles [11, Thm. 4.3.3], and it follows from the definition [21, Def. 1.7 (b), p. 36, (2), p. 37] that the rest are the points such that invariant sections of arbitrarily high powers of $\hat{\mathcal{L}}$ vanish; i.e., these form the base locus of high powers of $\hat{\mathcal{L}}$. Next, we claim that every unstable bundle appears in some $\hat{R}$. But any bundle $\mathcal{E}$ generated by its global sections fits into an extension of $\det \mathcal{E}$ by $\mathcal{O}_C^{\oplus r-1}$ [1, Thm. 2, p. 426]. Thus, for sufficiently high n , we find that all $\mathcal{E}$ either semistable or equal to any chosen unstable $\mathcal{E}_0$ will be contained in the family

$$0 \rightarrow \mathcal{O}(-n)^{\oplus r-1} \rightarrow \mathcal{E} \rightarrow (\det \mathcal{E}) \otimes \mathcal{O}(n(r-1)) \rightarrow 0,$$

simply by ensuring that n is high enough so that $\mathcal{E}(n)$ is globally generated. Fixing n , if we choose $n' \gg n$ high enough so that $\mathcal{E}(n')$ is globally generated and also $H^1(C, \mathcal{E}(n')) = 0$ for all $\mathcal{E}$ in the above Ext family, we find we can map this entire family into a corresponding GIT Quot scheme, with the semistable locus mapping into R . But since the family is irreducible, it maps into $\bar{R}$, and we see from the definition of $\hat{R}$ that in fact it maps into $\hat{R}$, as desired. Finally, we need to argue that the stack statement follows. We remark that any $\hat{R}$ has quotient stack isomorphic to an open substack of $\mathcal{M}_r(C)$ containing $\mathcal{M}_r^{ss}(C)$, and that furthermore as we vary our choices of $\hat{R}$, we obtain a cover of $\mathcal{M}_r(C)$. Also, using [2, Lem. 8.2] we see that G -invariant line bundles on $\hat{R}$ are in bijection with line bundles on $\mathcal{M}_r(C)$, so $\hat{\mathcal{L}}$ must be a power of the pullback of $\mathcal{L}$. Now, the sections of $\mathcal{L}^{\otimes n}$ on $\mathcal{M}_r(C)$ are in bijection with the sections on $\mathcal{M}_r^{ss}(C)$ [2, Prop. 8.3], which are in bijection with G -invariant sections of the pullback on $\bar{R}^{ss} = R$ [2, Lem. 7.2], which are in turn in bijection with the G -invariant sections of the pullback on $\hat{R}$ [23, Lem. 4.15]. Thus, every G -invariant section of the pullback of $\mathcal{L}$ on $\hat{R}$ is the pullback of a section on $\mathcal{M}_r(C)$, and since the varying choices of $\hat{R}$ cover $\mathcal{M}_r(C)$, the base locus of $\mathcal{L}^{\otimes n}$ on $\mathcal{M}_r(C)$ is the union of the (images of the) base loci on the different $\hat{R}$, which is precisely the unstable bundles, as desired.

For (iii), normality is evident because the GIT quotient uses rings of invariants of regular, hence integrally closed rings. Next, given (i), the argument of [18, 9.2] shows that $M_r(C)$ is $\mathbb{Q}$ -factorial. $\square$

We now finish the proof of the theorem.

Proof of Theorem A.6. We begin by proving part (i), by showing that V_r cannot be extended over the complement of U_r . The argument is in two parts. First, we show that if we know that the complement of U_r has codimension at least 2, then we obtain the desired result. We follow the notation of the preceding background theorem. Choose n sufficiently large, so that $\mathcal{L}^{\otimes n}$ and $(\mathcal{L}^{(p)})^{\otimes n}$ have base locus precisely equal to the locus of unstable bundles, and descend to very ample line bundles $\bar{\mathcal{L}}$ and $\bar{\mathcal{L}}^{(p)}$ on $M_r(C)$ and $M_r(C^{(p)})$. Let $(\bar{\mathcal{L}}, \bar{V})$ be the complete linear series for $\bar{\mathcal{L}}$ on $M_r(C)$. We wish to show that $V_r^*(\bar{\mathcal{L}}, \bar{V})$ extends uniquely to a

linear series $((\bar{\mathcal{L}}^{(p)})^{\otimes p}, V^{(p)})$ on $M_r(C^{(p)})$, whose base locus is precisely the locus of Frobenius-unstable bundles. The uniqueness of this extension would then imply that V_r cannot be extended over any of the Frobenius-unstable locus, as desired. Now, uniqueness follows from normality of $M_r(C^{(p)})$, together with the codimension hypothesis on the complement of U_r . To see existence, we pull back to $\mathcal{M}_r^{ss}(C)$, where by [2, Prop. 8.3], we can extend to a linear series $(\mathcal{L}^{\otimes n}, V)$ on all of $\mathcal{M}_r(C)$, which must have base locus precisely equal to the unstable locus. Using [17, Prop. A.1], we pull this back under the Verschiebung to $((\mathcal{L}^{(p)})^{\otimes np}, V^{(p)})$ on $\mathcal{M}_r(C^{(p)})$, which has base locus precisely equal to the locus of bundles whose Frobenius-pullback are unstable. Finally, by [2, Prop. 8.4] we can descend this to the desired $((\bar{\mathcal{L}}^{(p)})^{\otimes p}, V^{(p)})$.

To complete the proof of (i), we show that the complement of U_r has codimension at least 2. Now, let Z be any codimension 1 closed subscheme of $M_r(C^{(p)})$; we claim that it necessarily intersects any positive-dimensional closed subscheme of $M_r(C^{(p)})$. Indeed, using the fact that $M_r(C^{(p)})$ is $\mathbb{Q}$ -factorial, some multiple of Z is necessarily Cartier, and in particular there is an effective Cartier divisor with set-theoretic support equal to Z . Moreover, because $\text{Pic}(\mathcal{M}_r(C^{(p)})) \cong \mathbb{Z}$, this Cartier divisor is necessarily a descent to $M_r(C^{(p)})$ of some multiple of $\mathcal{L}^{(p)}$, which is ample, so must intersect any positive-dimensional closed subscheme, as claimed. But now we are done; the locus of direct sums of r line bundles inside $M_r(C^{(p)})$ gives a g^{r-1} -dimensional closed subscheme of $M_r(C^{(p)})$ which is visibly contained in U_r , so we conclude that the codimension of U_r is at least 2, as desired.

We now prove part (ii). We simply exhibit a point of $M_n(C^{(p)})$ at which V_n induces a finite flat map on versal deformation spaces. This point will correspond to a vector bundle $\mathcal{E}_0$ of the form $\mathcal{L}_1 \oplus \mathcal{L}_2 \oplus \cdots \oplus \mathcal{L}_n$, where the $\mathcal{L}_i$ are distinct line bundles of degree 0, with $\bigotimes_i \mathcal{L}_i \cong \mathcal{O}_{C^{(p)}}$, so that $\mathcal{E}_0$ has trivial determinant. By [29, p. 119], and in particular the asserted vanishing of $h^0(B \otimes L_1)$ for general L_1 and sequence (3), we may also require that the natural maps

$$H^1(C^{(p)}, \mathcal{L}_i \otimes \mathcal{L}_j^{-1}) \xrightarrow{F^*} H^1(C, F^* \mathcal{L}_i \otimes F^* \mathcal{L}_j)$$

are isomorphisms for all $i \neq j$.

We will consider deformations of $\mathcal{E}_0$ preserving the triviality of the determinant. One can check that this satisfies the criteria (H₁), (H₂), (H₃) of [30], so we let $R^{(p)}$ be a hull for the deformation problem and $\mathcal{E}$ on $C^{(p)} \times \text{Spec } R^{(p)}$ be the corresponding versal deformation of $\mathcal{E}_0$. The first-order deformations are parametrized by $H^1(C^{(p)}, \mathcal{E}nd^0(\mathcal{E}_0))$, and deformations are visibly unobstructed, so we conclude that $R^{(p)} \cong k[[t_1, \dots, t_N]]$, with $N = h^1(C^{(p)}, \mathcal{E}nd^0(\mathcal{E}_0))$. Since

$$\mathcal{E}nd^0(\mathcal{E}_0) \cong \mathcal{O}_{C^{(p)}}^{\oplus n-1} \oplus \bigoplus_{i \neq j} \mathcal{L}_i \otimes \mathcal{L}_j^{-1},$$

Riemann-Roch for vector bundles gives

$$\begin{aligned} h^1(C^{(p)}, \mathcal{E}nd^0(\mathcal{E}_0)) &= h^0(C^{(p)}, \mathcal{E}nd^0(\mathcal{E}_0)) - \deg \mathcal{E}nd^0(\mathcal{E}_0) + (\text{rk } \mathcal{E}nd^0(\mathcal{E}_0))(g-1) \\ &= (n-1) - 0 + (n^2-1)(g-1) = (g-1)n^2 + n - g. \end{aligned}$$

Now, let $I^{(p)} \subset R^{(p)}$ be the ideal defining the maximal closed subscheme of $\text{Spec } R^{(p)}$ over which $\mathcal{E}$ remains isomorphic to a direct sum of n distinct line bundles. If $J^{(p)}$ denotes the Jacobian of $C^{(p)}$, and $(J^{(p)})^n \rightarrow J^{(p)}$ is the addition morphism, let $T^{(p)}$ be the fiber of this morphism over 0. Since the kernel of the addition

map corresponds precisely to n -tuples of line bundles whose direct sum has trivial determinant, we may then describe $R^{(p)}/I^{(p)}$ as the completion of the local ring of $T^{(p)}$ at the point corresponding to $\mathcal{L}_1, \dots, \mathcal{L}_n$. Now, if we set

$$\mathcal{F}_0 := F^* \mathcal{E}_0 \cong \bigoplus_i F^* \mathcal{L}_i,$$

we can as before let R be a hull for the deformations of $\mathcal{F}_0$, and I the ideal cutting out the locus preserving the direct sum decomposition. We then obtain the corresponding description of R/I as above. Now, it is easily checked that the pullback under Frobenius induces a morphism $v : R \rightarrow R^{(p)}$. Next, since direct sum decompositions are preserved under pullback, $v(I) \subset I^{(p)}$, so we get an induced homomorphism $R/I \rightarrow R^{(p)}/I^{(p)}$. This clearly corresponds to the morphism $T^{(p)} \rightarrow T$ induced by the Verschiebung morphism $V : J^{(p)} \rightarrow J$; this last is finite flat, so we find that $R/I \rightarrow R^{(p)}/I^{(p)}$ is also finite flat.

Now, we also assert that we have

$$\mathrm{Hom}_k(I^{(p)}/\mathfrak{m}_{R^{(p)}} I^{(p)}, k) \cong \bigoplus_{i \neq j} H^1(C^{(p)}, \mathcal{L}_i \otimes \mathcal{L}_j^{-1})$$

and similarly for $\mathrm{Hom}_k(I/\mathfrak{m}_R I, k)$; we illustrate the argument for R . By the definition of a hull, $\mathfrak{m}_R/\mathfrak{m}_R^2$ is dual to the space of first-order infinitesimal deformations $\mathcal{T}_{\mathcal{F}_0}$ of $\mathcal{F}_0$, with the subspace $\mathcal{T}'_{\mathcal{F}_0} \subset \mathcal{T}_{\mathcal{F}_0}$ obtained by modding out by I and then dualizing corresponding to deformations preserving the direct sum decomposition. Thus, the quotient $\mathcal{T}_{\mathcal{F}_0}/\mathcal{T}'_{\mathcal{F}_0}$ corresponds to deformations transverse to those preserving the direct sum decomposition, and is obtained as the dual of the subspace $I/\mathfrak{m}_R I \subset \mathfrak{m}_R/\mathfrak{m}_R^2$. Now, it is also easy to see that $\mathcal{T}'_{\mathcal{F}_0}$ also corresponds to the summand of $H^1(C, \mathcal{E}nd^0(\mathcal{F}_0))$ given by $H^1(C, \mathcal{O}_C^{\oplus n-1})$. Thus, we see that $\mathcal{T}_{\mathcal{F}_0}/\mathcal{T}'_{\mathcal{F}_0}$ is given by $H^1(C, \bigoplus_{i \neq j} F^* \mathcal{L}_i \otimes F^* \mathcal{L}_j^{-1}) = \bigoplus_{i \neq j} H^1(C, F^* \mathcal{L}_i \otimes \mathcal{L}_j^{-1})$, which is the desired statement for R .

As a result, we find by our additional hypotheses on the choice of the $\mathcal{L}_i$ that v induces an isomorphism $I/\mathfrak{m}_R I \xrightarrow{\sim} I^{(p)}/\mathfrak{m}_{R^{(p)}} I^{(p)}$; we claim that this together with the finite-flatness of $R/I \rightarrow R^{(p)}/I^{(p)}$ is enough to imply that $v : R \rightarrow R^{(p)}$ is itself flat. Indeed, the isomorphism $I/\mathfrak{m}_R I \xrightarrow{\sim} I^{(p)}/\mathfrak{m}_{R^{(p)}} I^{(p)}$ implies that a set of generators of $I/\mathfrak{m}_R I$ over R will map to a set of generators of $I^{(p)}/\mathfrak{m}_{R^{(p)}} I^{(p)}$ over R and in particular over $R^{(p)}$, which by Nakayama's lemma implies that any lifts generate $I^{(p)}$ over $R^{(p)}$; we conclude that $I^{(p)} = v(I)R^{(p)}$. Now, we note that because $I_r \subset \mathfrak{m}_R$, we may write $R^{(p)}/v(\mathfrak{m}_R)R^{(p)} = (R^{(p)}/v(I)R^{(p)})/v(\mathfrak{m}_R)(R^{(p)}/v(I)R^{(p)}) = (R^{(p)}/I^{(p)})/v(\mathfrak{m}_R)(R^{(p)}/I^{(p)})$, which must be finite over R/I and hence over R because $R^{(p)}/I^{(p)}$ is. We thus have that the closed fiber of $\mathrm{Spec} R^{(p)} \rightarrow \mathrm{Spec} R$ is finite, and since both rings are regular of the same dimension, by [6, Thm 18.16 b] the map is flat, hence dominant. It remains to check that the map on hulls being dominant implies that the map on coarse moduli spaces is dominant. Because the coarse moduli spaces are irreducible (see [25, Rem. 5.5]), it suffices to show that the image of the map from the hull to the coarse moduli space is not contained in any proper closed subset, which follows easily from the definitions. $\square$

We conclude with a lemma on the case $g = 2, r = 2$ case which we used to conclude the Verschiebung is given by polynomials of degree p in Section 2.

Lemma A.9. *When $g = r = 2$, the line bundle $\mathcal{O}(1)$ on $M_2(C) \cong \mathbb{P}^3$ pulls back to the inverse of the determinant bundle on the stack $\mathcal{M}_2(C)$.*

Proof. We first use the criterion of [19, Prop. 2.2 (B)] to descend the determinant bundle $\mathcal{L}$ away from $\mathcal{O}_C \oplus \mathcal{O}_C$. On this locus, one easily check that the two conditions are satisfied for points with closed orbits, as these are the polystable bundles. Because $\mathbb{P}^3$ is regular, we can then extend uniquely to a descent of $\mathcal{L}$ on all of $M_2(C)$. Finally, since we already noted above that the inverse of the determinant bundle is the ample generator of $\text{Pic}(\mathcal{M}_r(C))$, the only bundle it could descend to is $\mathcal{O}(1)$. $\square$

APPENDIX B. A COMMUTATIVE ALGEBRA DIGRESSION

We develop some simple but non-standard commutative algebra over non-reduced rings which will be helpful in studying deformations of vector bundles. Throughout this section, R will denote a (typically non-reduced) Noetherian ring, $\mathfrak{n}$ the ideal of nilpotents of R , and M a finitely generated R -module, although we drop the Noetherian and finite-generated hypotheses for the first lemma. Note that everything we define in this section will be equivalent to their standard versions whenever R is integral.

Definition B.1. M is *NR-free* of rank r if M is generated by some $m_1, \dots, m_r$, such that given any relation $\sum_i a_i m_i = 0$, all the a_i must be nilpotent.

Lemma B.2. *If M is NR-free of rank r , then M_{red} is free of rank r . In particular, rank is well-defined for NR-free modules. Further, the converse holds if M is finitely generated and R is local, or if $\mathfrak{n}^m = 0$ for some m .*

Proof. Suppose M is NR-free of rank r . Clearly, the m_i still generate M_{red} , but using the fact that the m_i generate M , one checks that any non-zero relation modulo the nilpotents of R would yield a relation in R with not all coefficients nilpotent. Conversely, suppose M_{red} is free of rank r , with generators m_i . We show that any lifts of the m_i generate M , and they then clearly satisfy the desired relations restrictions. If R is local with maximal ideal $\mathfrak{m}$, and M finitely generated, this follows immediately from Nakayama's lemma. If, on the other hand, $\mathfrak{n}^m = 0$, we will get the desired result by showing via induction that $M' := M/(\{m_i\}_i)$ is contained in (the image of) $\mathfrak{n}^j M$ for all j . The base case is $j = 0$, which is a triviality. Now suppose it holds for $j - 1$, and we want to show it for j . Take $m \in M$; by the induction hypothesis, we can write $m = m' + \sum_i a_i m_i$ for some $m' \in \mathfrak{n}^{j-1} M$. Since the m_i generate M_{red} , writing out m' explicitly as a sum of products of elements of $\mathfrak{n}$, we find that there exist $a'_i \in \mathfrak{n}^{j-1}$ such that $m' - \sum_i a'_i m_i \in \mathfrak{n}^j M$, giving the desired result. $\square$

This argument also immediately gives us:

Corollary B.3. *Given two generating sets, m_i and m'_i for an NR-free R -module M , there is a (non-unique) invertible matrix T relating the m_i to the m'_i , such that if $\bar{m}_i$, $\bar{m}'_i$, and $\bar{T}$ are the images of m_i , m'_i and T in M_{red} , $\bar{T}$ is the matrix relating the $\bar{m}_i$ to the $\bar{m}'_i$.*

We can extend the standard definitions.

Definition B.4. M is **locally NR-free** of rank r if M becomes NR-free of rank r over every local ring of R . Given a separated Noetherian scheme X , a coherent sheaf $\mathcal{F}$ of $\mathcal{O}_X$ -modules is locally NR-free of rank r if $\mathcal{F}(U)$ is locally NR-free of rank r over $\mathcal{O}_X(U)$ for every affine U . Because our modules are finitely generated,

this is equivalent to there being an open cover of X on which $\mathcal{F}$ becomes NR -free of rank r .

We see that while we can attach transition matrices to locally NR -free sheaves, they are not unique, nor do they uniquely determine $\mathcal{F}$. However, they do determine $\mathcal{F}_{\text{red}}$ over X_{red} , which is all we will need for our purposes.

To develop the results we want, we will make the hypothesis for the rest of this section that $\text{Spec } R$ is irreducible. With this, we can prove the following proposition, which properly reformulated will kill several birds with one stone:

Proposition B.5. *Suppose that R_{red} is integral, and let M be a free R -module of rank r , and N an R -module generated by $n_1, \dots, n_s$, with ℓ non-zero relations of the form $fn_i = 0$ for some $f \in R$. Then if ϕ is a surjective map from M to N , $\ker \phi$ is an NR -free R -module of rank $r - s + \ell$. If further $\ell = 0$, then $\ker \phi$ is in fact free.*

Proof. First, we can suppose that f is not a unit, as if it were N would just be a free module of rank $s - \ell$, and we could simply use $n_{\ell+1}, \dots, n_s$ as generators. Our first claim is that if we lift ϕ from N to $\tilde{N} := R[n_1, \dots, n_s]$, the map will remain surjective. Let $\tilde{\phi}$ be some such lift, one can check by induction on j that $\tilde{N}/\text{im } \tilde{\phi} \subset f^j \tilde{N}/\text{im } \tilde{\phi}$ for all j . The Krull Intersection Theorem then implies that for some $f' \in (f)$, we have $(1 - f')(\cap_j (f^j)) = 0$. Now, by the hypothesis that R_{red} is integral, either $1 - f'$ is nilpotent, or $\cap_j (f^j) = 0$. However, if $1 - f'$ were nilpotent, f' would have to be a unit, which is not possible, since f was assumed not to be a unit. It follows that $\cap_j (f^j) = 0$, and $\tilde{\phi}$ is surjective.

Next, since it is a surjective map between free modules, the kernel of $\tilde{\phi}$ must be free, of rank $r - s$. This immediately proves the last assertion of the theorem. Now, let $\ker \tilde{\phi}$ be generated by $\tilde{m}_1, \dots, \tilde{m}_{r-s}$. Next, let $\hat{m}_1, \dots, \hat{m}_\ell$ be any elements mapping to $fn_1, \dots, fn_\ell \in \tilde{N}$ under $\tilde{\phi}$. It is then a routine check from the definitions $\{\tilde{m}_i, \hat{m}_j\}$ is a generating set making $\ker \phi$ into an NR -free module, of rank $r - s + \ell$. $\square$

Remark B.6. Note in particular that if f is a nonzero nilpotent, we will obtain modules of different rank by first taking the kernel and then restricting to R_{red} than by restricting to R_{red} and then taking the kernel.

To apply this proposition, we define:

Definition B.7. An **effective NR -Cartier divisor** on a Noetherian, separated, irreducible scheme X is a global section of the monoid sheaf $(\mathcal{O}_X \setminus \{0\})/\mathcal{O}_X^*$.

Lemma B.8. *Associated to any non-trivial effective NR -Cartier divisor f on X is a canonical closed immersion $X_f \hookrightarrow X$. Given any closed subscheme of X , there is at most one effective NR -Cartier divisor which induces it.*

Proof. Clear. $\square$

Remark B.9. Although X_f need not have codimension 1 in X for an effective NR -Cartier, it behaves in certain ways as if it had codimension 1, as the following theorem demonstrates.

Theorem B.10. *Let $X_f \hookrightarrow X$ be the closed immersion associated to a nontrivial effective NR -Cartier divisor f on X , $\mathcal{F}$ a locally free $\mathcal{O}_X$ -module of rank r on X , and $\mathcal{G}$ a coherent $\mathcal{O}_X$ -module on X . Let $f : \mathcal{F} \rightarrow \mathcal{G}$ be surjective. Then:*

- (i) if $\mathcal{G}$ is locally free of rank s on X , $\ker f$ is locally free of rank $r - s$, and
- (ii) if $\mathcal{G}$ is the pushforward of a locally free sheaf of rank s on X_f , $\ker f$ is locally NR -free on X of rank r .

Proof. (i) follows immediately from Proposition B.5, by restricting to a cover on which $\mathcal{F}$ and $\mathcal{G}$ are free, and noting that we are in the case $\ell = 0$. (ii) also follows from the same proposition, as if our cover is also fine enough give trivializing elements for f , we find that f matches up with the f of the proposition, and because $\mathcal{G}$ is a pushforward from X_f , it satisfies the hypotheses of the proposition with $\ell = s$, giving us the desired conclusion. $\square$

We will immediately apply this to prove a theorem to the effect that in certain cases, when one wants to replace an unstable vector bundle in a family with a semistable one, it suffices to look simply at the first order deformation induced by the family. We begin with the following lemma.

Lemma B.11. *Let $f : S \hookrightarrow T$ be a closed immersion of schemes, $\mathcal{F}$ a locally free sheaf on T , and $\mathcal{G}$ a locally free sheaf on S . Given $\psi : \mathcal{F} \rightarrow f_*\mathcal{G}$, then ψ arises via adjointness from $f^*\psi : f^*\mathcal{F} \rightarrow f^*f_*\mathcal{G} = \mathcal{G}$, and if $f^*\psi$ is surjective, then so is ψ . In this case, there is a natural surjection $f^*\ker\psi \twoheadrightarrow \ker f^*\psi$.*

Proof. Routine. $\square$

Theorem B.12. *Let $f : S \hookrightarrow T$ be a closed immersion, with T reduced, and such that if $i : S_{\text{red}} \hookrightarrow S$ is the standard inclusion, then both i and $f \circ i$ are induced by effective NR -Cartier divisors. Let $\mathcal{F}$ be a locally free sheaf of rank r on T , and $\mathcal{G}$ a locally free sheaf of rank s on S_{red} , and $\psi : \mathcal{F} \rightarrow f_*i_*\mathcal{G}$ arising via adjointness from a surjective map $f^*\mathcal{F} \rightarrow i_*\mathcal{G}$, which must then be $f^*\psi$. Then $i^*f^*\ker\psi = i^*\ker f^*\psi$, and both are locally free of rank r on S_{red} .*

Proof. The previous lemma gives us that ψ is surjective, and a surjection $f^*\ker\psi \twoheadrightarrow \ker f^*\psi$. Since pullback is right exact, this gives us $i^*f^*\ker\psi \twoheadrightarrow i^*\ker f^*\psi$, and by Theorem B.10 (i), it suffices to show these are both locally free of rank r , since then we would have the kernel of this surjection also being locally free, of rank 0. On the other hand, by (ii) of the same theorem, $\ker\psi$ is locally NR -free on T of rank r , hence locally free of rank r by reducedness of T . Thus, $i^*f^*\ker\psi$ must also be locally free of rank r . But likewise, since $f^*\mathcal{F}$ must also be locally free of rank r , using the same theorem, $\ker f^*\psi$ is locally NR -free of rank r on S , which as we saw initially is equivalent to $i^*\ker f^*\psi$ being locally free of rank r on S_{red} , as desired. $\square$

REFERENCES

- [1] M. F. Atiyah, *Vector bundles over an elliptic curve*, Proc. London Math. Soc. (3) **7** (1957), 414–452.
- [2] Arnaud Beauville and Yves Laszlo, *Conformal blocks and generalized theta functions*, Communications in Mathematical Physics **164** (1994), no. 2, 385–419.
- [3] Andrei Caldararu, *Derived categories of twisted sheaves on Calabi-Yau manifolds*, PhD thesis, Cornell University.
- [4] A. J. de Jong, *A conjecture on arithmetic fundamental groups*, Israel J. Math. **121** (2001), 61–84.
- [5] Pierre Deligne, *Théorie de Hodge. II*, Institut Des Hautes Etudes Scientifiques Publications Mathématiques (1971), no. 40, 5–57.
- [6] David Eisenbud, *Commutative algebra with a view toward algebraic geometry*, Springer-Verlag, 1995.

- [7] Gerd Faltings, *Algebraic loop groups and moduli spaces of bundles*, Journal of the European Mathematical Society **5** (2003), 41–68.
- [8] Eberhard Freitag and Reinhardt Kiehl, *Etale cohomology and the Weil conjecture*, Springer-Verlag, 1980.
- [9] Alexandre Grothendieck, *Le groupe de Brauer, II: Théorie cohomologique*.
- [10] ———, *Le groupe de Brauer, III: Exemples et compléments*, Dix exposés sur la cohomologie des schémas, Masson & Cie, 1968, pp. 88–188.
- [11] Daniel Huybrechts and Manfred Lehn, *The geometry of moduli spaces of sheaves*, Max-Planck-Institut für Mathematik, 1997.
- [12] Kirti Joshi and Eugene Z. Xia, *Moduli of vector bundles on curves in positive characteristic*, Compositio Math. **122** (2000), no. 3, 315–321.
- [13] Nicholas M. Katz, *Nilpotent connections and the monodromy theorem: Applications of a result of Turrittin*, Inst. Hautes Etudes Sci. Publ. Math. **39** (1970), 175–232.
- [14] Herbert Lange and Christian Pauly, *On Frobenius-destabilized rank-2 vector bundles over curves*, arXiv:math.AG/0309456.
- [15] Adrian Langer, *Semistable sheaves in positive characteristic*, Annals of Mathematics **159** (2004), no. 1, 251–276.
- [16] Yves Laszlo and Christian Pauly, *The action of the Frobenius maps on rank 2 vector bundles in characteristic 2*, Journal of Algebraic Geometry **11** (2002), no. 2, 129–143.
- [17] ———, *The Frobenius map, rank 2 vector bundles and Kummer’s quartic surface in characteristic 2 and 3*, Advances in Mathematics **185** (2004), no. 2, 246–269.
- [18] Yves Laszlo and Cristoph Sorger, *The line bundles of the moduli of parabolic g -bundles over curves and their sections*, Ann. Sci. École Norm. Sup. **30**, 499–525.
- [19] V. B. Mehta and T. R. Ramadas, *Moduli of vector bundles, Frobenius splitting, and invariant theory*, Annals of Mathematics **144** (1996), 269–313.
- [20] Shinichi Mochizuki, *Foundations of p -adic Teichmüller theory*, American Mathematical Society, 1999.
- [21] D. Mumford, J. Fogarty, and F. Kirwan, *Geometric invariant theory*, third enlarged ed., Springer-Verlag, 1994.
- [22] David Mumford, *Abelian varieties*, second ed., Oxford University Press, 1974.
- [23] M. S. Narasimhan and T. R. Ramadas, *Factorization of generalized theta functions. I*, Inventiones Mathematicae **114** (1993), no. 3, 565–653.
- [24] M. S. Narasimhan and S. Ramanan, *Moduli of vector bundles on a compact Riemann surface*, Annals of Mathematics **89** (1969), no. 2, 14–51.
- [25] P. E. Newstead, *Introduction to moduli problems and orbit spaces*, Springer-Verlag, 1978.
- [26] B. Osserman, *Frobenius-unstable bundles and p -curvature*, arXiv:math.AG/0409266.
- [27] ———, *Logarithmic connections with vanishing p -curvature*, arXiv:math.AG/0409145.
- [28] ———, *Mochizuki’s crys-stable bundles: A lexicon and applications*, arXiv:math.AG/0410323.
- [29] M. Raynaud, *Sections des fibrés vectoriels sur une courbe*, Bull. Soc. Math. France **110** (1982), no. 1, 103–125.
- [30] Michael Schlessinger, *Functors of Artin rings*, Transactions of the AMS **130** (1968), 208–222.